

# The complex Langevin equation and sign problem in lattice QCD

Dénes Sexty

Wuppertal University, Eötvös University

Budapest, 4<sup>th</sup> of May 2016

Collaborators: Gert Aarts, Erhard Seiler, Ion-Olimpiu Stamatescu  
Felipe Attanasio, Lorenzo Bongiovanni, Benjamin Jäger,  
Zoltán Fodor, Sándor Katz, Csaba Török

1. Sign problem (of lattice QCD)
  2. Ideas to solve it
  3. Some results extrapolating from  $\mu=0$
  4. Complex Langevin equation
    - Toy models
    - Lattice models – HDQCD and full QCD
- 
-

# Path integral formulation of QCD

Euclidean SU(3) gauge theory with fermions:

4d lattice    Temporal extent = inverse temperature

$$Z = \int DA_{\mu}^a D\bar{\Psi} D\Psi \exp(-S_E[A_{\mu}^a] - \bar{\Psi} D_E(A_{\mu}^a) \Psi)$$

$$A_{\mu}^a(x) \rightarrow U_{\mu}(x)$$

$$D_E(A) \rightarrow M(U) \text{ fermion matrix}$$

Integrating out fermions

$$Z = \int DU \exp(-S_E[U]) \det(M(U))$$

Haar measure of SU(3) group

SU(N) is compact



Finite volume of gauge orbits

No gauge fixing necessary

No Fadeev-Popov complication

We are interested in a system  
Described with the partition sum:

$$Z = \text{Tr} e^{-\beta(H - \mu N)} = \sum_C W[C]$$

Typically exponentially many configurations,  
no direct summation possible.

If the Weight is positive, build a Markov chain with the Metropolis alg.

$$\dots \rightarrow C_{i-1} \rightarrow C_i \rightarrow C_{i+1} \rightarrow \dots$$

Probability of visiting C

$$p(C) = \frac{1}{N_w} W[C]$$

$$\langle X \rangle = \frac{1}{Z} \text{Tr} X e^{-\beta(H - \mu N)} = \frac{1}{N_w} \sum_C W[C] X[C] = \frac{1}{N} \sum_i X[C_i]$$

This works if we have  $W[C] \geq 0$

Otherwise we have a **Sign problem**

# QCD sign problem

Euclidean SU(3) gauge theory with fermions:

$$Z = \int DU \exp(-S_E[U]) \det(M(U))$$


for  $\det(M(U)) > 0$  Importance sampling is possible  $\longrightarrow$  Hadron masses, EOS, ...

## Non-zero chemical potential

For nonzero chemical potential, the fermion determinant is complex

$$\det(M(U, -\mu^*)) = (\det(M(U), \mu))^*$$

Sign problem  $\longrightarrow$  Naive Monte-Carlo breaks down



# Sign problems in high energy physics

## Real-time evolution in QFT

“strongest” sign problem  $e^{iS_M}$

## Non-zero density (and fermionic systems)

$$Z = \text{Tr} e^{-\beta(H - \mu N)} = \int DU e^{-S[U]} \det(M[U])$$



Many systems: Bose gas  
XY model  
SU(3) spin model  
Random matrix theory  
QCD

## Theta therm

$$S = F_{\mu\nu} F^{\mu\nu} + i\Theta \epsilon^{\mu\nu\theta\rho} F_{\mu\nu} F_{\theta\rho}$$

And everything else with complex action

$$w[C] = e^{-S[C]} \quad w[C] \text{ is positive} \iff S[C] \text{ is real}$$



# How to solve the sign problem?

Probably no general solution - There are sign problems which are NP hard

[Troyer Wiese (2004)]

Many solutions for particular models with sign problem exist

Transforming the problem to one with positive weights

$$Z = \text{Tr} e^{-\beta(H - \mu N)} = \sum_C W[C] = \sum_D W'[D] \quad \begin{array}{l} \text{Dual variables} \\ \text{Worldlines} \end{array}$$

$$Z = \text{Tr} e^{-\beta(H - \mu N)} = \sum_n Z_n e^{\beta \mu n} \quad \text{Canonical ensemble}$$

$$Z = \text{Tr} e^{-\beta(H - \mu N)} = \int dE \rho_\mu(E) e^{-\beta E} \quad \text{Density of states}$$

$$Z = \text{Tr} e^{-\beta(H - \mu N)} = \sum_C W[C] = \sum_S \left( \sum_{C \in S} W[C] \right) \quad \text{Subsets}$$

## How to solve the sign problem?

### Extrapolation from a positive ensemble

Reweighting 
$$\langle X \rangle_W = \frac{\sum_c W_c X_c}{\sum_c W_c} = \frac{\sum_c W'_c (W_c/W'_c) X_c}{\sum_c W'_c (W_c/W'_c)} = \frac{\langle (W/W') X \rangle_{W'}}{\langle W/W' \rangle_{W'}}$$

Taylor expansion 
$$Z(\mu) = Z(\mu=0) + \frac{1}{2} \mu^2 \partial_\mu^2 Z(\mu=0) + \dots$$

Analytic continuation from imaginary sources  
(chemical potentials, theta angle,..)

### Using analyticity (for complexified variables)

Complex Langevin

Complexified variables – enlarged manifolds

Lefschetz thimble

Integration path shifted onto complex plane

In QCD direct simulation only possible at  $\mu=0$

Taylor extrapolation, Reweighting, continuation from imaginary  $\mu$ , canonical ens.  
all break down around

$$\frac{\mu_q}{T} \approx 1 - 1.5 \quad \frac{\mu_B}{T} \approx 3 - 4.5$$

Around the transition temperature

Breakdown at

$$\mu_q \approx 150 - 200 \text{ MeV}$$


$$\mu_B \approx 450 - 600 \text{ MeV}$$

Results on

$$N_T = 4, N_F = 4, ma = 0.05$$

using

Imaginary  $\mu$ ,  
Reweighting,  
Canonical ensemble




Agreement only at  $\mu/T < 1$




# Is there a critical point?

Critical point in  $(T, \mu)$  plane  $\rightarrow$   
 Critical surface in  $(T, \mu, m_{ud}, m_s)$  space




Order of the transition at  $\mu=0$



Which Scenario?

Calculate curvature at  $\mu=0$



[de Forcrand, Philipsen 2007...]

# Stochastic Quantization

Parisi, Wu (1981)

Given an action  $S(x)$

Stochastic process for  $x$ :

$$\frac{dx}{d\tau} = -\frac{\partial S}{\partial x} + \eta(\tau)$$

Gaussian noise  $\langle \eta(\tau) \rangle = 0$

$$\langle \eta(\tau) \eta(\tau') \rangle = \delta(\tau - \tau')$$

Averages are calculated along the trajectories:

$$\langle O \rangle = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T O(x(\tau)) d\tau = \frac{\int e^{-S(x)} O(x) dx}{\int e^{-S(x)} dx}$$

Fokker-Planck equation for the probability distribution of  $P(x)$ :

$$\frac{\partial P}{\partial \tau} = \frac{\partial}{\partial x} \left( \frac{\partial P}{\partial x} + P \frac{\partial S}{\partial x} \right) = -H_{FP} P$$

Real action  $\rightarrow$  positive eigenvalues

for real action the  
Langevin method is convergent

# Langevin method with complex action

The field is complexified

$$\frac{dx}{d\tau} = -\frac{\partial S}{\partial x} + \eta(\tau)$$

real scalar  $\longrightarrow$  complex scalar

link variables: SU(N)  $\longrightarrow$  SL(N,C)  
compact  $\longrightarrow$  non-compact

$$\det(U) = 1, \quad U^\dagger \neq U^{-1}$$

Analytically continued observables

$$\frac{1}{Z} \int P_{comp}(x) O(x) dx = \frac{1}{Z} \int P_{real}(x, y) O(x + iy) dx dy$$

$$\langle x^2 \rangle_{real} \rightarrow \langle x^2 - y^2 \rangle_{complexified}$$

“troubled past”: Lack of theoretical understanding  
Convergence to wrong results  
Runaway trajectories

Klauder '83, Parisi '83, Hueffel, Rumpf '83, Karsch, Wyld '84, Gausterer, Klauder '86.  
Matsui, Nakamura '86, ...

Interest went down as difficulties appeared

Renewed interest in connection of otherwise unsolvable problems

applied to nonequilibrium: Berges, Stamatescu '05, ...

aimed at nonzero density QCD: Aarts, Stamatescu '08 ... many important results since revival

# New results about complex Langevin in the last decade or so

1. Study many exactly solvable toy models to gain more understanding  
one-plaquette model, random matrices, thirring model,  
few variable models, Polyakov chain
2. Theoretical discussion and practical methods  
Proof of convergence  
Gauge cooling  
Non-holomorphic actions
3. Lattice models  
Non-equilibrium QFT (scalar field theory, pure gauge theory)  
XY model, SU(3) spin models, Bose gas  
Applications also in condensed matter: Bose gas in rotating frame,  
Imbalanced Fermi gas
4. Approximations to QCD  
HDQCD, kappa expansion
5. full QCD

# Proof of convergence

If there is fast decay  $P(x, y) \rightarrow 0$  as  $x, y \rightarrow \infty$

and a holomorphic action  $S(x)$

then CLE converges to the correct result

[Aarts, Seiler, Stamatescu (2009)

Aarts, James, Seiler, Stamatescu (2011)]

## (very rough) Sketch of the proof

$P(x, y, t)$ : probability density on the complex plane at Langevin time  $t$

$\rho(x, t)$ : complex measure evolving with the Fokker-Planck equation

$$\partial_t \rho(x, t) = \partial_x (\partial_x + (\partial_x S)) \rho(x, t)$$

$$\text{Stationary solution: } \rho(x, \infty) = \exp(-S(x))$$

CLE works, if  $\langle O(x) \rangle_{\rho(t)} = \langle O(x+iy) \rangle_{P(t)}$

$$\langle O(x) \rangle_{\rho(t)} = \langle O(x+iy) \rangle_{P(t)}$$

Interpolating function:

$$F(t, \tau) = \int P(x, y, t - \tau) O(x + iy, \tau) dx dy$$

$$F(t, 0) = \langle O(x + iy) \rangle_{P(t)} \quad F(t, t) = \dots = \langle O(x) \rangle_{\rho(t)}$$

$\partial_\tau F(t, \tau) = 0$  can be seen with partial integrations QED

## Non-holomorphic action for nonzero density

$$S = S_W[U_\mu] + \ln \text{Det } M(\mu)$$

measure has zeros ( $\text{Det } M = 0$ )  
 complex logarithm has a branch cut  
 —————▶ meromorphic drift

[Mollgaard, Splittorff (2013), Greensite(2014)]

is it a practical problem?

Depends on representation [Mollgaard, Splittorff (2014)]

Gauge cooling for eigenvals. [Nagata, Nishimura, Shimasaki (2016)]

Sufficient condition for correctness:

$$P(x, y) = 0 \quad \text{around the singularities}$$

[Aarts, Seiler, Sexty, Stamatescu in prep.]

# Gaussian Example

$$S[x] = \sigma x^2 + i\lambda x$$

CLE




$$\frac{d}{d\tau}(x+iy) = -2\sigma(x+iy) - i\lambda + \eta$$

$$P(x, y) = e^{-a(x-x_0)^2 - b(y-y_0)^2 - c(x-x_0)(y-y_0)}$$

Gaussian distribution  
around critical point

$$\left. \frac{\partial S(z)}{\partial z} \right|_{z_0} = 0$$

Measure  
on real axis



# Simple model of QCD with finite chemical potential

Euclidean U(1) One plaquette model with “fermion determinant”


$$Z = \int_0^{2\pi} dx e^{-S_B} \det M \quad S_B = -\frac{\beta}{2}(U + U^{-1}) = -\beta \cos(x) \quad U = e^{ix}$$

$$\det M = 1 + \frac{1}{2} \kappa (e^\mu U + e^{-\mu} U^{-1}) = 1 + \kappa \cos(x - i\mu)$$

Similar to QCD fermion determinant:

$$\det M(\mu) = [\det M(-\mu)]^* \quad \det M(i\mu) \quad \text{is real}$$

Exact averages calculated by numerical integration





# Fixedpoint structure



Distribution centered around attractive fixedpoints of the flow

$\mu$  grows



Fixed points move  
no change in analytical structure

No breakdown,  
Langevin works for high  $\mu$



# Gauge theories and CLE

## Stochastic quantisation on the group manifold

Updating must respect the group structure:

$$\langle \eta_{i_a} \rangle = 0$$

$$U'_i = \exp\left(i \lambda_a \left(-\epsilon D_{i,a} S[U] + \sqrt{\epsilon} \eta_{i,a}\right)\right) U_i$$

$$\langle \eta_{i_a} \eta_{j_b} \rangle = 2 \delta_{ij} \delta_{ab}$$

Left derivative: 
$$D_a f(U) = \left. \frac{\partial}{\partial \alpha} f(e^{i \lambda_a \alpha} U) \right|_{\alpha=0}$$

$\lambda_a$  Gellmann matrices

complexified link variables

$$\text{SU}(N) \longrightarrow \text{SL}(N, \mathbb{C}) \quad \det(U) = 1, \quad U^\dagger \neq U^{-1}$$

$$\text{compact} \longrightarrow \text{non-compact}$$

Distance from SU(N)

$$\sum_{ij} |(U U^\dagger - 1)_{ij}|^2$$

Unitarity Norms:

$$\text{Tr}(U U^\dagger) \geq N$$

$$\text{Tr}(U U^\dagger) + \text{Tr}(U^{-1} (U^{-1})^\dagger) \geq 2N$$

For SU(2):  $(\text{Im Tr } U)^2$

# Gauge cooling

[Seiler, Sexty, Stamatescu '13]

Minimize unitarity norm  
Distance from SU(N)  $\sum_i \text{Tr}(U_i U_i^\dagger - 1)$

Gauge transformation at  $x$  changes 2d link variables

$$U_\mu(x) \rightarrow \exp(-\alpha \epsilon \lambda_a G_a(x)) U_\mu(x)$$

$$U_\mu(x - a_\mu) \rightarrow U_\mu(x - a_\mu) \exp(\alpha \epsilon \lambda_a G_a(x))$$

Dynamical steps are interspersed  
with several gauge cooling steps


Gauge cooling leaves Fokker-Planck eq.  
For gauge invariant quantities unchanged  
[Nagata, Nishimura, Shimasaki '15]

Empirical observation:  
Cooling is effective for


$$\beta > \beta_{\min}$$

$$a < a_{\max}$$

Steepest descent



but remember,  $\beta \rightarrow \infty$   
in cont. limit  
 $a_{\max} \approx 0.1 - 0.2 \text{ fm}$




Smaller cooling

excursions into complexified manifold

“Skirt” develops  
small skirt gives correct result

propability distribution of Re Tr U



# Heavy Quark QCD at nonzero chemical potential (HDQCD)

Hopping parameter expansion of the fermion determinant  
Spatial hoppings are dropped

$$\text{Det } M(\mu) = \prod_x \det(1 + C P_x)^2 \det(1 + C' P_x^{-1})^2$$

$$P_x = \prod_\tau U_0(x + \tau a_0) \quad C = [2\kappa \exp(\mu)]^{N_\tau} \quad C' = [2\kappa \exp(-\mu)]^{N_\tau}$$

$$S = S_W[U_\mu] + \ln \text{Det } M(\mu)$$


Studied with reweighting


[De Pietri, Feo, Seiler, Stamatescu '07]

$$R = e^{\sum_x C \text{Tr } P_x + C' \text{Tr } P^{-1}}$$

CLE study using gaugecooling

[Seiler, Sexty, Stamatescu (2012)]






Gauge cooling stabilizes the distribution  
 SU(3) manifold instable even at  $\mu=0$

Fermion density:

$$n = \frac{1}{N_\tau} \frac{\partial \ln Z}{\partial \mu}$$

average phase:

$$\langle \exp(2i\varphi) \rangle = \left\langle \frac{\text{Det } M(\mu)}{\text{Det } M(-\mu)} \right\rangle$$




$$\det(1 + CP) = 1 + C^3 + C \text{Tr } P + C^2 \text{Tr } P^{-1}$$


Sign problem is absent at  
small or large  $\mu$

Reweighting is impossible at  $6 \leq \mu/T \leq 12$ , CLE works all the way to saturation

# Comparison to reweighting



$6^4$  lattice,  $\beta=5.9$



$6^4$  lattice,  $\mu=0.85$




Discrepancy of plaquettes at  $\beta \leq 5.6$   
 a skirted distribution develops

$$a(\beta=5.6) = 0.2 \text{ fm}$$



# Mapping the phase diagram

[Aarts, Attanasio, Jäger, Seiler, Sexty, Stamatescu, in prep.]



fixed  $\beta=5.8 \rightarrow a \approx 0.15 \text{ fm}$

$\kappa=0.12$


onset transition at  $\mu = -\ln(2\kappa) = 1.43$

$N_t * 8^3$  lattice


$N_t = 2..28$

Temperature scanning

# Exploring the phase diagram of HDQCD




Onset in fermionic density  
Silver blaze phenomenon



Polyakov loop  
Transition to deconfined state

$$\beta=5.8 \quad \kappa=0.12 \quad N_f=2 \quad N_t=2\dots 24$$

# Polyakov loop susceptibility



Hint of first order deconfinement and first order onset transition

# Extension to full QCD with light quarks [Sexty (2014)]

QCD with fermions  $Z = \int DU e^{-S_G} \det M$

Additional drift term from determinant

$$K_{axv}^F = \frac{N_F}{4} D_{axv} \ln \det M = \frac{N_F}{4} \text{Tr} (M^{-1} M'_{va}(x, y, z))$$

Noisy estimator with one noise vector

Main cost of the simulation: CG inversion

Inversion cost highly dependent on chemical potential

Eigenvalues not bounded from below by the mass

(similarly to isospin chemical potential theory)


Unimproved staggered and Wilson fermions

Heavy quarks: compare to HDQCD

Light quarks: compare to reweighting

# CLE and full QCD with light quarks [Sexty (2014)]

Physically reasonable results



Non-holomorphic action  
poles in the fermionic drift  
Is it a problem for full QCD?

So far (at high temperatures), it isn't:  
Comparison with reweighting  
Study of the spectrum  
Hopping parameter expansion



# Reweighting

$$\langle F \rangle_{\mu} = \frac{\int DU e^{-S_E} \det M(\mu) F}{\int DU e^{-S_E} \det M(\mu)} = \frac{\int DU e^{-S_E} R \frac{\det M(\mu)}{R} F}{\int DU e^{-S_E} R \frac{\det M(\mu)}{R}}$$
$$= \frac{\langle F \det M(\mu) / R \rangle_R}{\langle \det M(\mu) / R \rangle_R} \quad R = \det M(\mu=0), |\det M(\mu)|, \text{ etc.}$$

$$\left\langle \frac{\det M(\mu)}{R} \right\rangle_R = \frac{Z(\mu)}{Z_R} = \exp\left(-\frac{V}{T} \Delta f(\mu, T)\right)$$

$\Delta f(\mu, T)$  = free energy difference

Exponentially small as the volume increases  $\langle F \rangle_{\mu} \rightarrow 0/0$

Reweighting works for large temperatures and small volumes




Sign problem gets hard at  $\mu/T \approx 1$

# Comparison with reweighting for full QCD


[Fodor, Katz, Sexty, Török 2015]

Reweighting from ensemble at

$$R = \text{Det } M(\mu=0)$$



# Overlap problem



Histogram of weights  
Relative to the largest weight in ensemble



Average becomes dominated by very few configurations



# Sign problem

Sign problem gets hard around

$$\mu/T \approx 1 - 1.5$$



$$\langle \exp(2i\phi) \rangle = \left\langle \frac{\det M(\mu)}{\det M(-\mu)} \right\rangle$$




# Comparisons as a function of beta

Similarly to HDQCD

Cooling breaks down at small beta


at  $N_T=4$  breakdown at  $\beta=5.1 - 5.2$

At larger  $N_T$  ?




# Comparisons as a function of beta

$N_T=6$



$N_T=8$



Breakdown prevents simulations in the confined phase  
for staggered fermions with  $N_T=4,6,8$

Two ensembles:  $m_\pi \approx 4.8 T_c$   
 $m_\pi \approx 2.3 T_c$

# Conclusions

Sign problem of lattice QCD      solid results only below  $\mu_q/T=1$

Evading the sign problem by direct simulations  
using complexified fields in the **Complex Langevin Equation**

Recent progress for CLE simulations

Better theoretical understanding (poles?)

Gauge cooling

Phase diagram of HDQCD mapped out


full QCD with light quarks – only high temperatures so far

Outstanding issues

- What happens if the poles are problematic?  
How to diagnose, how to solve the problem?
- is QCD at low temperatures an example for that?

# Spectrum of the Dirac Operator $N_F=4$ staggered

Massless staggered operator at  $\mu=0$  is antihermitian



# Spectrum of the Dirac Operator

$N_F=4$  staggered

